

TEG Kaolin (with and without Heparinase)



Alternate Name:	TEG test
Performing Lab:	New Hanover
Specimen Container:	Blue stopper tube (sodium citrate)
Minimum Volume Required:	Tube must be filled completely
Testing Availability	Routine: 24 hours/day Stat: Yes
Special Handling:	None
Patient Preparation:	None
Specimen Stability:	2 hours
Reference Range:	Reaction Time: 5-10 minutes Clot Kinetics: 1-3 minutes Clot Angle: 53-72 deg Maximum Amplitude: 50-70 mm Lysis after 30 minutes:0-8%
Critical Value:	None
CPT Code:	85347,85384x2,85576
Testing Methodology:	Thrombelastograph
Causes for Specimen Rejection:	Improper labeling Clotted specimen Specimen diluted with IV fluid or contaminated with heparin Specimens that are too old Under or over-filled collection tube
Clinical Significance:	The TEG® system monitors shear elasticity (a physical property), it is sensitive to all the interacting cellular and plasmatic components in the blood that may affect the rate or structure of a clotting sample and its breakdown. The overall profile can be qualitatively or quantitatively interpreted in terms of the hypo, normal or hypercoagulable state of the sample, the degree of lysis, and other measurements of coagulopathy.